

D-instantons on orbifolds and gauge/gravity correspondence

Alessandro Tanzini ¹

¹ LPTHE, Université Paris VI, and INFN, Roma.

e-mail: tanzini@lpthe.jussieu.fr

Abstract: D-instantons are used to probe the near-horizon geometry of D3-branes systems on orbifold spaces. For fractional D3-branes, D-instanton calculus correctly reproduces the gauge β -function and $U(1)_R$ anomaly of the corresponding $\mathcal{N} = 2$ non-conformal Super Yang-Mills theories. For D3-branes wrapping the orbifold singularity, D-instantons can be identified with gauge instantons on ALE space, providing evidence of AdS/CFT duality for gauge theories on curved spaces.

The duality between $\mathcal{N} = 4$ Super Yang-Mills theory (SYM) in four dimensions and type IIB string theory compactified on $AdS_5 \times S^5$ [1] has provided a first concrete example of how a string theory can be extracted from a gauge theory. A remarkable result is that the multi-instanton moduli space of $SU(N)$ $\mathcal{N} = 4$ SYM reproduces in the large N limit the ten dimensional near-horizon $AdS_5 \times S^5$ geometry of the dual type IIB supergravity [2]. This allows to establish a careful correspondence between instanton dominated correlators in gauge theory and D-instanton corrections to the supergravity action, thus providing a non trivial check of the AdS/CFT duality.

Recently, many interesting generalizations of this duality to more realistic non-conformal gauge theories have been considered. In this context, the study of type IIB string theory on orbifold spaces play a major rôle. A general class of models with a lower amount of supersymmetry is in fact defined by locating a stack of D3-branes at a $\mathbb{R}^6/\Gamma$ singularity, where Γ is a discrete subgroup of $SO(6) \cong SU(4)_R$. Depending on whether Γ is embedded in an $SU(4)$, $SU(3)$ or $SU(2)$ subgroup of $SU(4)_R$, the resulting low-energy Yang-Mills theories display respectively $\mathcal{N} = 0, 1, 2$ supersymmetry [3]. Moreover, if one consider fractional branes, which do not have images under the action of the group Γ , the corresponding gauge theories turn out to be non-conformal [4]; various supergravity solutions of this kind have been studied for the $\mathcal{N} = 2$ case [5]. However, in all these models the short distance region of the space-time is excised by an enhançon mechanism, and additional states have to be taken into account to describe this region. In the gauge theory, the enhançon radius corresponds to the scale where instanton effects become relevant, thus one can expect that D-instantons come into play on the supergravity side. In [6], the low-energy dynamics of the D(-1)-D3 system in presence of fractional branes have been considered, along the lines of the analysis carried out in [2, 7] for the conformal case. The D-instanton partition function has been shown to factorize *already at finite* N in a term associated to the overall center of mass degrees of freedom, which describes the geometry of the space where the D-instantons are moving, and an effective matrix

model describing internal degeneracies of the k D-instantons bound state. In the present discussion, we focus on the pure $\mathcal{N} = 2$ SYM theory. A more general analysis can be found in [6], to which we refer also for the notations and conventions and for a complete list of references. The pure $\mathcal{N} = 2$ SYM action describes the low-energy dynamics of a stack of N fractional D3-branes lying at a $\mathbb{R}^4/\Gamma$ singularity [5]. This theory can be obtained by means of a Γ -projection on the $\mathcal{N} = 4$ $SU(N)$ gauge theory associated to N nearby D3-branes moving on a flat space-time. Analogously, the multi-instanton moduli space measure and multi-instanton action of pure $\mathcal{N} = 2$ can be read from the low-energy lagrangian describing the dynamics of a fractional D(-1)-D3 system on $\mathbb{R}^4/\Gamma$. We recall that the effective theory for k D(-1) and N D3-branes moving on flat space is given by the dimensional reduction to $0 + 0$ dimensions of an $\mathcal{N} = 1$ $U(k)$ gauge theory in $D = 6$ with vector multiplet $(\chi_a, \lambda_{\dot{\alpha}A}, D_c)$, one hypermultiplet transforming in the adjoint representation $(a_{\alpha\dot{\alpha}}, \mathcal{M}_\alpha^A)$ and N transforming in the fundamental representation (and its conjugate), denoted by $(w_{\dot{\alpha}}, \mu^A; \bar{w}^{\dot{\alpha}}, \bar{\mu}^A)$ ¹. The low-energy excitations of the fractional D(-1)-D3 system are defined by the Γ -projection on these fields, which for $\Gamma = \mathbb{Z}_p$ reads²

$$\begin{aligned}\mu^A &= e^{2\pi i \frac{q_A}{p}} \mu^A & \bar{\mu}^A &= e^{2\pi i \frac{q_A}{p}} \bar{\mu}^A \\ \mathcal{M}_\alpha^A &= e^{2\pi i \frac{q_A}{p}} \mathcal{M}_\alpha^A & \lambda_{\dot{\alpha}}^A &= e^{2\pi i \frac{q_A}{p}} \lambda_{\dot{\alpha}}^A \\ \chi^{AB} &= e^{2\pi i \frac{q_A + q_B}{p}} \chi^{AB}\end{aligned}\tag{1}$$

with $q_1 = q_2 = 0$, $q_3 = -q_4 = 1$. All the other fields $(w_{\dot{\alpha}}, \bar{w}^{\dot{\alpha}}, a_m, D^c)$ are invariant under the $\mathbb{Z}_p$ orbifold action. The net effect of the projection (1) on the bosonic fields is to set to zero the components $(\chi_1, \chi_2, \chi_4, \chi_5)$ of the vector field χ_a , which roughly describes the position of the D-instantons in the transverse space to the D3-branes. Thus the fractional D-instantons are allowed to move only on the orbifold plane (χ_3, χ_6) . As for the fermionic fields, it is easily realized that the projection (1) breaks the $SU(4)$ $\mathcal{R}$ -symmetry group of the $\mathcal{N} = 4$ theory down to $SU(2)_{\dot{A}} \times U(1)_{\mathcal{R}}$, where $SU(2)_{\dot{A}}$ is the automorphism group and $U(1)_{\mathcal{R}}$ the anomalous $\mathcal{R}$ charge of the $\mathcal{N} = 2$ SYM theory. By introducing the complex coordinates on the orbifold plane

$$\begin{aligned}\phi &\equiv 2\chi_{43} = 2\chi^{12} = \frac{1}{\sqrt{2}}(-\chi_3 + i\chi_6) \\ \bar{\phi} &\equiv 2\chi_{21} = 2\chi^{34} = \frac{1}{\sqrt{2}}(-\chi_3 - i\chi_6) \ ,\end{aligned}\tag{2}$$

we can now write the $\mathcal{N} = 2$ multi-instanton action as

$$S_{k,N} = \frac{1}{g_0^2} S_G + S_K + S_D\tag{3}$$

with

$$\begin{aligned}S_G &= \text{tr}_k \left(2[\phi, \bar{\phi}]^2 + \frac{i\pi}{\sqrt{2}} \lambda_{\dot{\alpha}\dot{A}} [\phi, \lambda^{\dot{\alpha}\dot{A}}] - D^c D^c \right) \\ S_K &= \text{tr}_k \left(2\bar{\phi} L \phi - \sqrt{2} i \pi \bar{\phi} (\bar{\mu}_{\dot{A}} \mu^{\dot{A}} + \mathcal{M}_{\dot{A}}^\alpha \mathcal{M}_\alpha^{\dot{A}}) \right) \\ S_D &= \text{tr}_k \left(i\pi (-[a_{\alpha\dot{\alpha}}, \mathcal{M}^{\alpha\dot{A}}] + \bar{\mu}^{\dot{A}} w_{\dot{\alpha}} + \bar{w}^{\dot{\alpha}} \mu^{\dot{A}}) \lambda_{\dot{A}}^{\dot{\alpha}} + D^c (\bar{w} \tau^c w - i \bar{\eta}_{mn}^c [a_m, a_n]) \right) \ ,\end{aligned}\tag{4}$$

¹Here $a = 1, \dots, 6$ stands for the vector index in the $SO(6)$ group acting on the transverse space to the D3-branes, $\alpha, \dot{\alpha} = 1, 2$ for spinorial indices of the Lorentz group acting on the longitudinal space, and $A = 1, \dots, 4$ for the fundamental index of the internal $SU(4)_R \cong SO(6)$ group. Finally, $c = 1, 2, 3$ labels the auxiliary fields of the vector multiplet [6].

²We adopt the notations of [7]. Given the local group isomorphism $SO(6) \cong SU(4)$, $SO(6)$ vectors can also be written as $\chi_{AB} \equiv \Sigma_{AB}^a \chi_a / \sqrt{8}$, with Σ_{AB}^a given in terms of the t'Hooft symbols.

where we have defined the linear operator $L \cdot \Omega = \{W^0, \Omega\}/2 + [a_m, [a_m, \Omega]]$, with $W_{ij}^0 = \bar{w}_{iu}^\alpha w_{uj\dot{\alpha}}$. It is worth to remark that the multi-instanton action (3) can be easily twisted into a topological matrix model [6]. Thus, the parameters appearing in this action can be continuously deformed without affecting the results on the computation of the observables, as long as they are well defined. In particular, the partition function can be evaluated in the limit $g_0 \rightarrow \infty$; life is much simpler in this limit, since the term S_G in (3) is suppressed and the fields (D_c, λ_α^A) become lagrangian multipliers implementing the ADHM constraints in S_D [2]. In order to explicitly evaluate the partition function it is moreover convenient to define the $SU(N)$ gauge invariant variables $W_{ij}^m, \zeta_{ij}^{\dot{A}}$ through [2]

$$\begin{aligned} W_{ij}^0 &= \bar{w}_{iu}^\alpha w_{uj\dot{\alpha}} , & W_{ij}^c &= \bar{w}_{iu}^\alpha \tau_\alpha^{c\dot{\beta}} w_{uj\dot{\beta}} \quad c = 1, 2, 3 \\ \mu_{iu}^A &= w_{uj\dot{\alpha}} \zeta_{ji}^{\dot{A}} + \nu_{iu}^A, & \text{with } \bar{w}_{iu}^\alpha \nu_{uj}^A &= 0, \\ \bar{\mu}_{iu}^A &= \bar{\zeta}_{\dot{\alpha}ij}^A \bar{w}_{uj}^\alpha + \bar{\nu}_{iu}^A & \text{with } \bar{\nu}_{iu}^A w_{uj\dot{\alpha}} &= 0 \end{aligned} \quad (5)$$

and perform the integrations over the iso-orientation modes (parameterizing a point in the coset space $\frac{SU(N)}{SU(N-2k) \times U(1)}$) and over their “fermionic superpartners” $\nu^A, \bar{\nu}^A$. Then by integrating out the auxiliary field D^c we enforce the bosonic ADHM constraint $W^c = i\bar{\eta}_{mn}^c [a_n, a_m]$, while the integration on λ_α^A enforces the fermionic constraint. After these manipulations, we are finally left with the $SU(N)$ invariant measure [7]

$$\begin{aligned} Z_{k,N} &= c_{k,N} e^{2\pi i k \tau} \frac{1}{\text{Vol } U(k)} \int d^{k^2} W^0 d^{4k^2} a d^{k^2} \phi d^{k^2} \bar{\phi} d^{4k^2} \mathcal{M} d^{4k^2} \zeta \\ &\times \left(\det_{2k} W \det_k \bar{\phi}^2 \right)^{N-2k} e^{-S_{k,N}} , \end{aligned} \quad (6)$$

with

$$\begin{aligned} S_{k,N} &= -4\pi i \text{tr}_k \bar{\phi} \Lambda^{12} + \text{tr}_k (2\bar{\phi} L \phi) \\ \Lambda^{\dot{A}\dot{B}} &= \frac{1}{2\sqrt{2}} \left(\zeta^{[\dot{A}} W \zeta^{\dot{B}]} + [a^{\alpha\dot{\alpha}}, \mathcal{M}_\alpha^{[\dot{A}} \zeta_{\dot{\alpha}}^{\dot{B}]} + \mathcal{M}^{\alpha[\dot{A}} \mathcal{M}_\alpha^{\dot{B}]} \right) \\ c_{k,N} &= \frac{2^{2kN-4k^2+k} \pi^{2kN-2k^2+k}}{\prod_{i=1}^{2k} (N-i)!} (-2\pi^2)^{k(N-2k)} \pi^{4k^2} . \end{aligned} \quad (7)$$

The determinant factors in (6) come from the Jacobian of the change of variables (5) and from the integration on the fermionic variables $\nu^A, \bar{\nu}^A$.

We would like now to factor out the dependence on the center of mass degrees of freedom in (6) in order to get informations on the near-horizon geometry of our model. As we will see shortly, this can be done already at finite N . Inspired by large N manipulations of [2] we split each $U(k)$ adjoint V-field into its trace and traceless part

$$V = v_0 \mathbf{1}_{k \times k} + \hat{v} . \quad (8)$$

At this point it is convenient to rescale the various traceless components of the fields in the following way

$$\begin{aligned} a_m &= -x_m + \rho \hat{a}_m \\ \phi &= r e^{2i\vartheta} (1 + \hat{\phi}) \\ \bar{\phi} &= r e^{-2i\vartheta} (1 + \hat{\bar{\phi}}) \end{aligned}$$

$$\begin{aligned}
W_0 &= \rho^2 (1 + \hat{W}_0) \\
\zeta_{\hat{\alpha}}^{\dot{A}} &= e^{i\vartheta} \bar{\eta}_{\hat{\alpha}}^{\dot{A}} + \rho^{-\frac{1}{2}} e^{i\vartheta} \hat{\zeta}_{\hat{\alpha}}^{\dot{A}} \\
\mathcal{M}_{\alpha}^{\dot{A}} &= e^{i\vartheta} \xi_{\alpha}^{\dot{A}} + \rho \hat{a}_{\alpha\hat{\alpha}} e^{i\vartheta} \bar{\eta}^{\dot{A}\hat{\alpha}} + \rho^{\frac{1}{2}} e^{i\vartheta} \hat{\mathcal{M}}_{\alpha}^{\dot{A}} .
\end{aligned} \tag{9}$$

From (9) it follows that the variables $x_m, \rho^2, \xi_{\hat{\alpha}}^{\dot{A}}, \bar{\eta}_{\hat{\alpha}}^{\dot{A}}$ are the trace components associated to the overall bosonic and fermionic translational/conformal zero modes, while r is the distance of the D-instanton probe from the D3-branes³. Our results can be expressed in terms of this quantity and of the adimensional variable $y = r\rho$. In fact, plugging (9) in (6) and taking care of the Jacobian involved in the rescalings, one can easily check that all dependence in the center of mass variables r, x_m, ϑ factors out as

$$\begin{aligned}
Z_{k,N} &= d_{k,N} \int d^4 x_m r^3 dr d\vartheta e^{-4ikN\vartheta} r^{-2kN} e^{2\pi i k \tau_0} d^4 \xi d^4 \bar{\eta} \\
&= d_{k,N} \int d^4 x_m r^3 dr d\vartheta e^{2\pi i k \tau(r)} d^4 \xi d^4 \bar{\eta} .
\end{aligned} \tag{10}$$

In (10) we can easily recognize the $\text{AdS}_5 \times \text{S}^1$ volume form⁴, $d^4 x_m r^3 dr d\vartheta$, multiplied by a power of the complex coordinate $r e^{2i\vartheta}$. Remarkably, this deformation of the $\text{AdS}_5 \times \text{S}^1$ factor can be completely reabsorbed in the following redefinition of the complex coupling

$$2\pi i \tau(r, \vartheta) \equiv i\theta - \frac{8\pi^2}{g(r)^2} = i(\theta_0 - 4N\vartheta) - \frac{8\pi^2}{g_0^2} - 2N \ln(r) . \tag{11}$$

If, as usual in AdS/CFT correspondence, we identify the AdS radial coordinate r with the dynamical energy scale μ of the SYM gauge theory, we can read from the term multiplying the logarithm in (11) the first coefficient in the expansion of the β function for the gauge coupling

$$\beta \equiv r \frac{d}{dr} g(r) = -b_1 g^3 / 16\pi^2 + \dots . \tag{12}$$

with $b_1 = 2N$, in agreement with the field theory result. Moreover, the extra phase factor $e^{-4ikN\vartheta}$ in (10) correctly reproduces the $U(1)_R$ anomaly of $\mathcal{N} = 2$ SYM. Notice in fact that the angular rescalings in (9) correspond to an $U(1)_R$ transformation on the fields of the $\mathcal{N} = 2$ SYM. Due to the presence of instantons, such a transformation is not a symmetry of the lagrangian, but induces a variation of the θ_0 parameter precisely equal to that we have found in (11). This also imply that in order to have a non-trivial correlation function one should consider the insertion of suitable operators carrying $n = 4kN$ fermionic zero-modes. An interesting example could be the single or multi-trace composite operators of scalar fields, which have been shown not to get perturbative anomalous dimensions in $\mathcal{N} = 2$ theories [8]. Finally, the $d_{k,N}$ in (10) are numerical coefficients given by integrals on the left-over $SU(k)$ variables, which describe the internal dynamics of the k D-instanton bound state. They can be computed only in a limited number of cases, and notably in the large N limit where they are an important ingredient to check the AdS/CFT duality [2]. The formulation in terms of topological matrix theory is a crucial tool in this evaluation, and could be useful also for finite N computations. However, we remark that the coefficients $d_{k,N}$ do not involve powers of r , and therefore cannot affect the result found in (10).

³Actually $r = d/\alpha'$, where d is the distance from the D3-branes in the orbifold fixed plane.

⁴We use coordinates [1] where the $\text{AdS}_5 \times \text{S}^1$ metric reads $ds^2 = \frac{r^2}{R^2} dx_m^2 + \frac{R^2}{r^2} dr^2 + R^2 d\vartheta^2$ with $R^2 = \sqrt{4\pi g_s N}$, g_s being the string coupling constant.

Summarizing, the near-horizon geometry turns out to be a deformed $\text{AdS}_5 \times \text{S}^1$, where the deformation is entirely determined in terms of the one-loop β function and $U(1)_R$ anomaly of the underlying gauge theory. By means of the technique just described one can study also more general configurations containing regular D3-branes and regular D-instantons. Since these last possess images under the discrete group Γ , they can leave the orbifold plane and roam in the entire transverse space, probing the complete $\text{AdS}_5 \times \text{S}^5 / \mathbb{Z}_p$ geometry [6].

Let us now comment on the case in which the orbifold quotient is taken along the directions longitudinal to the D3-branes system [9]. This means that the discrete group Γ is now embedded in an $SU(2)$ factor of the Lorentz group $SO(4) \cong SU(2)_L \times SU(2)_R$, say for definiteness $SU(2)_L$. The projection is thus similar to that in (1), but now Γ is acting on the spinor indices α, β of $SU(2)_L$. Moreover, the orbifold group acts non trivially also on the Chan-Paton indices. Thus, starting from a solution with instanton number $K = \sum_q k_q$ and gauge group $U(N)$, $N = \sum_q N_q$, the surviving components turn out to be [6]

$$\begin{aligned} w_{\dot{\alpha}}^q &= w_{\dot{\alpha} i_q u_q} & \mu_q^A &= \mu_{i_q u_q}^A \\ a_q^{\alpha \dot{\alpha}} &= a_{i_q + q_{\alpha} j_q}^{\alpha \dot{\alpha}} & \mathcal{M}_q^{A\alpha} &= \mathcal{M}_{i_q + q_{\alpha} j_q}^{A\alpha} \\ \chi_q^{AB} &= \chi_{i_q j_q}^{AB} & D_q^c &= D_{i_q j_q}^c & \lambda_q^{A\dot{\alpha}} &= \lambda_{i_q j_q}^{A\dot{\alpha}} \end{aligned} \quad (13)$$

They correspond to block matrices with block indices $i_q = 1, \dots, k_q$, $u_q = 1, \dots, N_q$, and $q = 0, \dots, p-1$ for $\Gamma = \mathbb{Z}_p$. All sums in q are from now on understood modulo p . Notice that, contrary to the previous case (1), the whole $\mathcal{N} = 4$ supersymmetry is preserved by this projection since Γ acts in the same way on the different components of a given supermultiplet. The moduli space of multi-instanton solutions and ADHM constraints can be described in terms of the invariant components (13). In particular, the moduli space is spanned by the super-coordinates in the first two lines of (13). The presence of $\dim D^c = 3k_q^2$ bosonic ADHM constraints for each q and of the auxiliary gauge group $\prod_q U(k_q)$ reduces the number of independent bosonic degrees of freedom to

$$\dim \mathcal{M}_B = 4 \sum_q (k_q N_q + \hat{k}_q k_q - k_q^2) \quad , \quad (14)$$

where $\hat{k}_q = \frac{1}{2} \sum_{\alpha} k_{q+q_{\alpha}}$, with $q_1 = -q_2 = 1$. Analogously, due to the presence of $\dim \lambda^{A\dot{\alpha}} = 8k_q^2$ fermionic ADHM constraints for each q , the number of independent fermionic degrees of freedom is

$$\dim \mathcal{M}_F = 8 \sum_q (k_q N_q + \hat{k}_q k_q - k_q^2) \quad . \quad (15)$$

The resulting dimensions of the moduli space in (14) and (15) are in agreement with that obtained from the Kronheimer-Nakajima construction of gauge instantons on ALE spaces [10]. Thus, in presence of D3-branes wrapping an $\mathbb{R}^4 / \mathbb{Z}_p$ singularity, we are led, following [9], to identify D-instantons with self-dual connections for non-abelian gauge theories living on ALE spaces. An important new feature emerges in this case; indeed, since ALE manifolds support non trivial two-cycles, the first Chern class is in general non-vanishing and characterizes the instanton solution together with the second Chern class usually considered on flat spaces. In particular, instanton solutions with vanishing first Chern class must obey the condition [6]

$$N_q + 2k_q - 2\hat{k}_q = 0 \quad \text{for } q > 0. \quad (16)$$

We remark that only in these cases the second Chern class coincides with the instanton number defined as $K/|\Gamma| = K/p$. (16) is a highly non trivial constraint on the allowed values of (k_q, N_q) . In the simplest case, *i.e.* the $SU(2)$ bundle on the Eguchi–Hanson blown-down space $\mathbb{R}^4/\mathbb{Z}_2$, solutions to (16) are given by either $\vec{N} = (2, 0), \vec{k} = (k, k)$ or $\vec{N} = (0, 2), \vec{k} = (k-1, k)$. The corresponding instanton solutions have respectively integer and half-integer second Chern class, while the dimension of the moduli space computed from (14) turns out to be equal to $8k$ and $8k-4$ respectively, in agreement with [10]. A computation of the partition function for the lowest value of the Chern class, $c_2 = 1/2$, was carried out in [10] yielding the bulk contribution to the Euler number of the moduli space⁵.

The low energy action for this D(-1)–D3 system can be easily obtained by implementing the Γ -projection (13) on the $U(k)$ effective gauge theory of the parent D(-1)–D3 system living on flat space. Since the whole $SO(6)$ $\mathcal{R}$ -symmetry is clearly preserved by the projection, this action describes instantons in an $\mathcal{N} = 4$ gauge theory, living on an ALE space. It is a straightforward exercise to apply the techniques previously described to show that for a regular D-instanton probe the center of mass degrees of freedom factor out from the multi-instanton measure and describe a point in an euclidean $S^5 \times AdS_5^E/\mathbb{Z}_p$ space, with $\mathbb{Z}_p$ acting on the four-dimensional $\mathbb{R}^4$ boundary of the AdS_5^E [6].

References

- [1] J. Maldacena, Adv. Theor. Math. Phys. **2** (1998) 231, [hep-th/9711200].
- [2] N. Dorey, T.J. Hollowood, V.V. Khoze, M.P. Mattis and S. Vandoren, Nucl. Phys. **B552** (1999) 88, [hep-th/9901128].
- [3] S. Kachru and E. Silverstein, Phys. Rev. Lett. **80** (1998) 4955, [hep-th/9802183]; A. Lawrence, N. Nekrasov and C. Vafa, Nucl. Phys. **B533** (1998) 199, [hep-th/9803015]; M. Bershadsky, Z. Kakushadze and C. Vafa, Nucl. Phys. **B523** (1998) 59, [hep-th/9803076].
- [4] I.R. Klebanov and N. Nekrasov, Nucl. Phys. **B574** (2000) 263, [hep-th/9911096].
- [5] for a review, see M. Bertolini, P. Di Vecchia and R. Marotta, hep-th/0112195.
- [6] F. Fucito, J.F. Morales and A. Tanzini, JHEP **0107** (2001) 012, [hep-th/0106061].
- [7] T.J. Hollowood, V.V. Khoze and M.P. Mattis, JHEP **9910** (1999) 019, [hep-th/9905209]; T. J. Hollowood and V.V. Khoze, Nucl.Phys. **B575** (2000) 78, [hep-th/9908035].
- [8] N. Maggiore and A. Tanzini, Nucl. Phys. **B613** (2001) 34, [hep-th/0105005].
- [9] M.R. Douglas and G. Moore, hep-th/9603167; C.V. Johnson and R.C. Myers, Phys. Rev. **D55** (1997) 6382, [hep-th/9610140].
- [10] M. Bianchi, F. Fucito, G. Rossi and M. Martellini, Phys. Lett. **B359** (1995) 56, [hep-th/9507003]; Nucl. Phys. **B473** (1996) 367, [hep-th/9601162].

⁵It is known and rigorously proven that for $c_2 = 1/2$ the moduli space is a copy of the base manifold (the Eguchi–Hanson manifold) whose bulk contribution is $3/2$.